



CERTIFICATE Of Appreciation



This Certificate is Awarded to

Oktavina Kartika Putri, M.Si., M.Sc.

Has Participated as

POSTER SESSIONS

at

**The 1st International Conference on Science Technology and Engineering
for Sustainable Development (ICOSTES) 2018**

September 10th-11st, 2018 at University of Islam Malang, East Java, Indonesia



Prof. Dr. Maskuri, M.Si.



Dr. Ir. Nurhidayati, M.P.



Comparison of Total Flavonoid Content on Fresh and Dried Fig (*Ficus carica*) Leaf with Various Brewing Temperature

by Oktavina Kartika Putri, M.Si., M.Sc.

oktavina.chemistry@gmail.com

ABSTRACT

Fig (*Ficus carica*) has the various benefits especially for health because of its various active compounds. One of the active substances found in fig is flavonoid. The use of fig leaves in the form of teabags certainly involves the heating process when brewing. Whereas, thermal processes have a great influence on the availability of flavonoid from foods. This study aims to compare the total flavonoid content on fresh and dried fig (*Ficus carica*) leaf with various brewing temperature. The temperatures used for brewing vary from 60°C, 70°C, and 80°C. The total flavonoid content was determined by spectrophotometric method with the standard quercetin solution. The total flavonoid contents of fresh fig leaf brew with the brewing temperatures of 60°C, 70°C, and 80°C are 0.0031±0.00039%, 0.0081±0.00021%, and 0.0092±0.00024%. The total flavonoid contents of dried fig leaf brew with the brewing temperatures of 60°C, 70°C, and 80°C are 0.0010±0.00025%, 0.0015±0.00004%, and 0.0017±0.00018%. Based on the Two Way Anova test, it can be concluded that the brewing temperature and simplicia preparation of fig leaf (fresh and dried) significantly affect the total flavonoid content. The fig leaf brew with the highest flavonoid content is 0.0092±0.00024% obtained from brewing the fresh fig leaf at 80°C.

Keywords: total flavonoid content, fresh fig leaf, dried fig leaf, brewing temperature

INTRODUCTION




OBJECTIVES


This study was conducted to compare the total flavonoid content on fresh and dried fig (*Ficus carica*) leaf with various brewing temperature.

RESULTS AND DISCUSSION

Fresh Fig Leaf Brew



Dried Fig Leaf Brew



The increase in temperature increases the total flavonoid content of dried fig leaf brew. This means that the maximum temperature used (80°C) is a safe temperature for the flavonoid extraction process. Based on the comparison of results, it can be seen that brew with the highest total flavonoid content is from fresh fig leaf with a brewing temperature of 80°C. In addition, with the same brewing temperature, total flavonoid content of fresh fig leaf brew was higher than dried one. The destruction of flavonoids had varied significant differences between fresh samples and dried materials. Heating may breakdown some phytochemicals which affect cell wall integrity and cause a migration of some flavonoids component.

METHODOLOGY



The main tool used for the total flavonoid content determination was the UV-Vis spectrophotometer (Hitachi U-2900).

Fresh and dried fig leaves were brewed with the temperature of 60°C, 70°C and 80°C.

CONCLUSION

Increasing the brewing temperature from 60°C to 80°C increases the total flavonoid content of fig leaf brew, both fresh and dried. Total phenolic content of fresh fig leaf brew is higher than dried fig leaf brew, because the drying process decreases the availability of flavonoids.

Thanks to:



PROCEEDING



PROCEEDING

The 1st International Conference
on Science Technology and
Engineering for Sustainable
Development

“Towards Enhancement
of Agriculture and Animal
Husbandry, Natural Resources
and the Environment,
and Engineering in the ASEAN”

September 10 -11- 2018
Malang, Indonesia

Held on Cooperation between :

University of Islam Malang
(Faculty of Agriculture, Animal
Husbandry, Mathematic and
Science, and Engineering) and
some abroad Universities.

Sponsored by :



PETROSIDA GRESIK
Always Valuable



PROCEEDING

International Conference on Science Technology and Engineering
For Sustainable Development
(ICoSTES) 2018

*“Towards Enhancement of Agriculture and Animal Husbandry Natural
Resources and the Environment and Engineering in the ASEAN”*

Malang, September 10-11th 2018



Publisher:

Universitas Islam Malang

Executive Committee

Chairman	: Nur Hidayati, Dr., Ir., MP
Vice Chairman	: 1. Inggit Kentjonowaty, Dr., Ir., MP 2. Tintrim Rahayu, Ir., M.Si 3. Unung Lesmanah, Ir., MT
Secretary	: 1. Anis Sholihah, Dr., Ir., MP 2. Ismi Choirotin, ST., MT., M.Sc
Finance treasurer	: 1. Sri Hindarti, Ir., Msi 2. Sri Susilowati, Ir., MM
IT & Website	: 1. Hasan Zayadi, SSi., MSi 2. Anis Rosyidah, Dr., Ir., MP
Event Division	: 1. Cynthia Permata Dewi, ST., MT., M.Sc. 2. Siti Asmaniyah M, SP., MP
Publication & Funding	: 1. Ratna Djuniwati, Dr., M.Si 2. Djuhari, Ir., MSi
General Accomodation	: 1. Irawati Dinasari, Ir., MP 2. Farida Syakir, Ir., MP 3. Ari Kurniawan, ST 4. Risna Amalia 5. Moch. Muzaqqi, SP 6. Puguh Darmawan, ST 7. Syamsudin

Steering Committee

Maskuri, Dr., MSi (University of Islam Malang)
Inocencio E Buot Jr, Ph. D (University of the Philippines Los Banos, Philippines)
Sutiman Bambang Sumitro, M.Sc., D.Sc (University of Brawijaya)

Reviewer

Inocencio E Buot Jr, Ph. D (University of the Philippines Los Banos, Philippines)
Nurhidayati, Dr., Ir., MP (University of Islam Malang)
Mudawamah, Dr., Ir., MSi (University of Islam Malang)
Merites M Buot, Ph. D (University of the Philippines Los Baños, University of the Philippines Open University, Philippines)
Nour Athiroh A.S, Dr., MKes., SSi (University of Islam Malang)
Novi Arfarita, SP, MP, M.Sc, Ph.D (University of Islam Malang)
M. Jasa Afroni, ST.,MT.,Ph.D (University of Islam Malang)
Ari Ambarwati, Dr., M.Pd (University of Islam Malang)
Nikmatul Khoiriyah, Ir., MP (University of Islam Malang)
Cynthia Permata Dewi, ST, MT, M.Sc (University of Islam Malang)
Ari Hayati Dr., Dra., MP (University of Islam Malang)
Ratna Djuniwati Lisminingsih, Dr., M.Si (University of Islam Malang)

Managing Editor:

Hasan Zayadi, S.Si, M.Si
Ismi Choirotin, ST., MT., M.Sc.

Held in Between:

Faculty of Agriculture, University of Islam Malang
Faculty of Mathematics and Science, University of Islam Malang

Faculty of Husbandry, University of Islam Malang
Faculty of Engineering, University of Islam Malang
University of the Philippines Los Banos, Philippines
National Central University, Taiwan
Brawijaya University, Indonesia
Yangon University, Myanmar
Suranaree University of Technology, Thailand
Kumamoto University, Japan

Publisher:

Universitas Islam Malang (University of Islam Malang)
Jl. Mayjend Haryono 193 Malang, 65144
Telp. 0341 551932
Fax. 0341 552249
Email: icostes2018@unisma.ac.id
ISBN: 978- 602-52411-3-0

1st Printing Edition, September 2019

No part of this publication may be reproduce without either prior written permission of the publisher

Table of Content

Editorial Board -- iv

Foreward -- v

Speech of Rector -- vii

Conferences Schedule -- ix

Keynote Speech

- Biodiversity in the ASEAN: Status, Challenges and Opportunities for Scientists and Universities
Inocencio E Buot Jr, PhD -- xi
- Application of clean energy with examples of solar energy
Jiunn-Chi Wu, PhD -- xii
- Biology as Inspiration in settling Community Problems
Sutiman Bambang Sumitro, M.Sc., D.Sc -- xiii
- Varieties of Myanmar Mangoes and Off-season Fruit Production of *Mangifera indica* L. cv. Sein ta lone
Thanda Aye, Ph.D. -- xv
- Sustainable Animal Production in the Tropics
Pramote Paengkoum, Ph.D -- xvi
- Climate Change Mitigation & Adaptation: Global to ASEAN & Malaysian Perspective (Special Focus on Agricultural Productivity)
Dr. Rawshan Ara Begum, Ph.D -- xvii
- Climate Change and Its Impact on Agricultural Sectors in Indonesia
Dr. Ir. Agus Sugianto, ST., MP. -- xviii

Table of Content -- xix

Sub Theme: Basic Science

- 1 Quercetin Compounds of Avocado Leaf Extracts (*Persea Americana, Mill*) as A Reducer of Total-Cholesterol Level
Elly Rustanti, Enny Puspita -- 1
- 2 Dipping Bikang (Seed and Peel of Black Grape) Potency in Reducing Milk Bacteria Contamination of Dairy Cattle
Dyanovita Al Kurnia, Qabillah Cita -- 7
- 3 Enhancement of Contents of Sunflower Oil with Abiotic Elicitor by Light Wavelength of Colours on *Calluses Helianthus annuus*
Dwie Retna Suryaningsih, Sri Arijanti Prakoeswa -- 12

- 4 Epinasty response on petiole angle of *Capsicum frutescens* L. in flooding stress
Muhammad Rizza Pahlevi, Serafinah Indriyani, Retno Mastuti, Estri Laras Arumingtyas -- 20
- 5 Utilization of ¹⁵N Labeled Rice Straw for Tracking Nitrogen Distribution with Indicator of Maize Plant
Anis Sholihah and Agus Sugianto -- 27
- 6 The influenced of cultivar red *Moringaoleifera* fermentation on number PMN cells of Liver histopathology mice (*Musmusculus*) after infected by Salmonella typhimurium
MM Riyaniarti Estri Wuryandaria, Widodo, Edi Widjanto, Muhaimin Rifa'i -- 35
- 7 Selection of Boer-Eb Lambs from Cross Breeding Based on Performance of Phenotype as A Superior Parental of Broiler Goat Types
Inggit Kentjnowaty, Farid Wadjdi, Sri Susilowati, Jaya Wulandari and Dodi Jarwoko -- 40
- 8 Study of Sub Chronic Administration with *Scurrula atropurpurea* to Blood Parameters
Nour Athiroh AS -- 45
- 9 Comparison of Total Flavonoid Content on Fresh and Dried Fig (*Ficus carica*) Leaf with Various Brewing Temperature
Oktavina Kartika Putri -- 49

Sub Theme: Applied Science

- 10 Molecular Weight Profile of The Chicken feet Antioxidant Peptides
Edy Susanto, Djalal Rosyidi, Lilik Eka Radiati, Subandi -- 53
- 11 Intensity of Shoot Fly Attacks (*Atherigona* sp) Due to The Late Planting of Sumenep Local Corn Talango Variety
Ida Ekawati, H. D. Wati, and Isdiantoni -- 59
- 12 Influence of Encapsulation Material on The Quality of Probiotic Containing *Lactobacillus salivarius*
Umi Kalsum, MF Wadjdi -- 63
- 13 Genetic Evaluation of Boer Sire Based on Performance of Its Crosses with Local Goats
Ratna Kumala Dewi, Wardoyo -- 67
- 14 The Application of Pesticides for Controlling Pest and Its Impact to Agriculture Ecosystem and Public Health
Luh Titi Handayani, Achmadi Susilo -- 72
- 15 The Giant Freshwater Prawn (*Macrobrachium rosenbergii de Maen*) with Closed System Transportation
Anny Rimalia, Yulius Kisworo -- 77
- 16 Effectiveness of Leave Soursoup Extract (*Annona Muricata*) and Leave Neem Extract (*Azadirachta Indica*) as Herbal Insecticide In Cattle Infected By Blood Sucker Fly (*Haematobia Exigua*)
Totok Dwi Ambodo, Muhammad Faiz Zidni Mubarak, Hasan Hilmi Aziz, Nurul Humaidah -- 82
- 17 Screening of Probiotic Bacteria Candidates From Digestive Tract of Tidal Swamp Fishes
Rini Marlida, Elrifadah -- 85
- 18 Innovation of Fermented Feeding Based on Waste Sludge Coconut as Low Cholesterol Duck Feeding
Fitriyah, Afidhatul Masruroh, Faiqul Mubarak, Nurul Humaidah -- 92
- 19 Attack Intensity of *Plutella xylostella* Pests and Yield of Cabbage (*Brassica oleraceae* L. var. *Capitata*) Due to Application of Vermicompost with Additive of Papaya Leaf and Its Mixture
Abdul Basit, Nurhidayati, Hesti Wahyu Lestari -- 96

- 20 The Effect of Sound Wave and Drought Stress on Stomatal Opening and Soybean Growth
Istirochah Pujiwati, Bambang Guritno, Nurul Aini, Setyawan, P. Sakti -- 102
- 21 Study of Bintaro Plants (Cerbera manghas) as Bioinsecticides Against Armyworm in Soybean Plants
Achmadi Susilo, Dwi Harjanta, Tatuk Tojibatus S -- 114

Sub Theme: Engineering

- 22 Expansive Clay Soil Stabilization Using Composite Portland Cement and Lime on The Road Subgrade
Frans Pratama, Suryo Hapsoro Tri Utomo, Imam Muthohar -- 120
- 23 A Drainage System for Inundation Problems At Subdistricts of Sidoarjo Regency
Anita Rahmawati -- 132
- 24 Automotive Industrial Electricity Demand Prediction Due to Lightweight Body to Support Multi-Purpose Electric Vehicle (MPEV) Massive Production
Gigih Adicita Wijaya, Rudy Setiabudi -- 136
- 25 Improvement of Farmer Harvesting Results by the Optimization Model of Microcontroller Based Irrigation
Bambang Suprpto, Tri Budi Prayogo, Ismi Choirotin, Eko Noerhayati -- 147
- 26 Unusual Ecg (Electrocardiogram) Pattern Detection Using Sax (Symbolic Aggregate Approximation) Algorithm
Mohammad Jasa Afroni, Bambang Minto -- 153
- 27 Model Sprinkler Irrigation Based on Gravity
Eko Noerhayati, Anita Rahmawati, Margianto, Bambang Dwisulo -- 160
- 28 BOD and COD Reduction Using Stratified Filter and Constructed Wetland
Lies Kurniawati Wulandari, M. Bisri, Donny Harisuseno, Emma Yuliani -- 165

Sub Theme: Social Science

- 29 The Use of Balanced Scorecard in Micro Small Medium Enterprises (MSME)
Siska Elvani, Jani Januar, Triana Dewi H -- 173
- 30 Development of Interactive Learning Media Based on Edutainment of Writing Skill Procedure Text
Kustyarini, Sri Utami, Endang Koesmijati -- 179
- 31 The Effect of Think, Pair and Share Learning Models and Attitudes on Cooperative Skills of Seventh Grade Students
Khoirul Efendiy -- 185
- 32 Multimodal Efficacy of Counseling in Students Stresses Management
Esy Suraeni Y, Kustyarini -- 192
- 33 Roles and Benefits of Mudharabah Contract Business Capital on Domestic Violence Woman in An-Nisa' Cooperative Pati Central Java
Rini Rahayu Kurniati, Daris Zunaida, Srinuring Wahyu -- 199
- 34 Analysis of Marketing Mix on Customer Satisfaction in Traditional Market
Budi Prihatminingtyas, Putri Budi Setyowati -- 206
- 35 Competitive Sustainability Tourism Strategy with Transglobal Leadership Approach to SME Tourism Sector in East Java
Adya Hermawati, Suhermin, Rahayu Puji Suci -- 211

- 36 Using SI GEMBUL Media in Improving Creative Thinking Ability
Istiqomah Aminin -- 217
- 37 Social Change in coffe farmers at Jember District
Isti Nurrohimah, Sri Subekti, Lenny Widjyanthi -- 221
- 38 Model of Human Resource Management Integrated Based Leadership in Strategic Efficiency Achieving Global Competitiveness
Rahayu Puji Suci, Adya Hermawati, Suwarta -- 229
- 39 Forest Farmers as Main Actor of Social Forestry Program
Indah Novita Dewi -- 236
- 40 A Review on The Different Responses of Arabica Coffee Farmers Due to Climate Change
Cinder Rosa, Sugeng Rahartoa, Luh Putu -- 245
- 41 Marketing Challenges of Fermented Cassava (Tape) Agroindustry in Jember
Khusna Ismiya Yanuasari, Triana Dewi, Lenny Widjyanthi -- 251
- 42 Performance Measurement The “An” Batik SMEs in Batu Using the Smart System Method
C F Putri, I Nugroho, D Purnomo -- 256
- 43 The Application of Digital Technology to Increase Agricultural Production in Indonesia
Dyanasari -- 265
- 44 Sangko’ Development as Diverse Alternative of Food Consumed by Poor Household in Marginal Area
Mubarokah, Syarif Imam Hidayat – 270
- 45 THE IMPACT OF PRICE CHANGES ON ANIMAL FOOD DEMAND IN EAST JAVA, INDONESIA: The Quadratic Almost Ideal Demand System Approach
Nikmatul Khoiriyah, Ratya Anindita, Nuhfil Hanani, Abdul Wahib Muhaimin --274
- List of Participant -- 284**

COMPARISON OF TOTAL FLAVONOID CONTENT ON FRESH AND DRIED FIG (*FICUS CARICA*) LEAF WITH VARIOUS BREWING TEMPERATURE

Oktavina Kartika Putri^a

^aAkademi Farmasi Putra Indonesia, Malang, Indonesia

*Corresponding author e-mail: oktavina.chemistry@gmail.com

ABSTRACT

Fig (*Ficus carica*) has the various benefits especially for health because of its various active compounds. One of the active substances found in fig is flavonoid. The use of fig leaves in the form of teabags certainly involves the heating process when brewing. Whereas, thermal processes have a great influence on the availability of flavonoid from foods. This study aims to compare the total flavonoid content on fresh and dried fig (*Ficus carica*) leaf with various brewing temperature. The temperatures used for brewing vary from 60°C, 70°C, and 80°C. The total flavonoid content was determined by spectrophotometric method with the standard quercetin solution. The total flavonoid contents of fresh fig leaf brew with the brewing temperatures of 60°C, 70°C, and 80°C are 0.0031±0.00039%, 0.0081±0.00021%, and 0.0092±0.00024%. The total flavonoid contents of dried fig leaf brew with the brewing temperatures of 60 °C, 70° C, and 80°C are 0.0010±0.00025%, 0.0015±0.00004%, and 0.0017±0.00018%. Based on the Two Way Anova test, it can be concluded that the brewing temperature and simplicia preparation of fig leaf (fresh and dried) significantly affect the total flavonoid content. The fig leaf brew with the highest flavonoid content is 0.0092±0.00024% obtained from brewing the fresh fig leaf at 80°C.

Keywords: total flavonoid content, fresh fig leaf, dried fig leaf, brewing temperature

1. Introduction

Tin or commonly called fig (*Ficus carica*) has the various benefits especially for health because of its various active compounds. One of the active substances found in fig is flavonoid. Flavonoids have antioxidant, antiviral and antibacterial properties (Koz³owska and Dorota, 2014). They also regulate gene expression and modulate enzymatic action (Pollastri and Tattini, 2011).

Compared to fruit, the flavonoid content of fig leaf is higher. Flavonoid levels of ethanolic extract of fig leaves and fruits respectively 2.62±0.003 and 1.96±0.002 mg rutin/g (Trifunschi *et al.*, 2015). Therefore, the researcher prefer to study the flavonoid content of fig leaf than its fruit. The main types of flavonoid found in fig leaves are quercetin and luteolin with amounts 631 dan 681 mg/kg fig leaves extracts (Vaya and Mahmood, 2006). Quercetin is the most important flavonoid which belongs to the class of flavonol which widely used in medicine and pharmaceuticals. Quercetin provided many health promoting benefits, like cardiovascular properties, cancer reducing agent, anti-inflammatory, asthma and many more (Kumar *et al.*, 2017). Therefore, determination of flavonoid contents in fig leaf brew performed with using standard quercetin solution.

The traditional and current uses of fig leaves are to treat cough, prevention of nutritional anaemia, anthelmintic, irritant potential, and tuberculosis (Mawa *et al.*, 2013). The use of fig leaves in Indonesia is starting to widely increase. Moreover, some industries have produced dried fig leaves that are packaged in the form of teabags. The use of fig leaves in the form of teabags certainly involves the heating process when brewing. Whereas, thermal processes have a great influence on the availability of flavonoid from foods. In fact, quercetin which is one of the flavonoid content in fig leaves is labile on heating (Kumar *et al.*, 2017).

This study was conducted to compare the total flavonoid content on fresh and dried fig (*Ficus carica*) leaf with various brewing temperature. The temperatures used for brewing vary from 60°C, 70°C, and 80°C. The use of dried and fresh fig leaves was to prove whether the drying process also affects flavonoid contents.

2. Material and Methods

2.1 Material

The fig leaf used are obtained from Blimbing, Lowokwaru, Malang and tested for its authenticity in Materia Medica Batu, East Java with the determination number of 074/253A/102.7/2018. The varieties, ages, parts of leaves, and harvest time of foliage are not determined. The main tool used for the total flavonoid content determination is the UV-Vis spectrophotometer (Hitachi U-2900). The other materials used were methanol, AlCl_3 2% and quercetin as a standard solution.

2.2 Methods

Fresh and Dried Fig Leaf Preparation

Fresh fig leaves are washed and then air dried and was cut with the width of about 2-3 mm. Thereafter, ± 6.25 grams the fresh fig leaf was put into 3 beaker glasses of each to be brewed. And then, ± 6.25 grams the fresh fig leaf was put into 3 evaporation plates of each to be dried with the temperature of 25-30°C with an oven until a constant weight is obtained. After that, the dried fig leaf moved into 3 beaker glasses to be brewed.

Fig Leaf Brewing

Aquadest was boiled, then cooled to a temperature of 60°C, 70 °C and 80 °C. After that, 100 mL aquadest was used to brew fresh and dried fig leaves of each. The brew obtained was closed until reaching a room temperature, then filtered.

Total Flavonoid Content Determination

Flavonoid contents of fresh and dried fig brew were determined by the AlCl_3 Colorimetric Method (Chandra *et al.*, 2014) with the standard quercetin solution. The first step in determining the flavonoid levels was to determine the calibration curve of standard quercetin solution. After got the curve equation, then the sample's absorbance was plotted in the equation. In brief to determine the absorbance of the sample, fig leaf brew was added with 4 mL of methanol then added with 1 mL of AlCl_3 2%. After that, the mixture was incubated at room temperature for 30 minutes. After incubation, the absorbance was measured at a maximum wavelength of 430 nm.

Data Analysis

The data obtained were analyzed with Two Way Anova test to find out:

- whether brewing temperature affects total flavonoid content or not,
- whether simplicia (fresh and dried) preparation affects total flavonoid content or not,
- whether the interactions of both (the brewing temperature and simplicia preparation) affect total flavonoid content or not.

3. Result and Discussion

Quercetin standard curves have been obtained with the curve equations $y = 0.079x - 0.0063$ and $R^2 = 0.9986$. After the absorbance of fresh fig leaf brew was plotted in the equation, the total flavonoid content was obtained and it is presented in Figure 3.1.




Figure 3.1 Total flavonoid content of fresh fig leaf brew

Based on the results, the largest total flavonoid content of fresh fig leaf brew obtained was from 80°C. Analyzed from the resulting diagram, from 60°C to 80°C the temperature increases the total flavonoid content. One of several factor that influences extraction speed is temperature. High temperature can increase the plant active compounds desorption due to the increase of cell damage to the material caused by the high solvent temperature (Jain *et al.*, 2009).

Total flavonoid content of dried fig leaf brew also showed the same trend as fresh fig leaf brew, it is presented in Figure 3.2. The increase in temperature increases the total flavonoid content of dried fig leaf brew. This is not in line with the statement which states that thermal processes have a great influence on the availability of flavonoid from foods (Kumar *et al.*, 2017). This means that the maximum temperature used (80°C) is a safe temperature for the flavonoid extraction process.




Figure 3.2 Total flavonoid content of dried fig leaf brew

Based on the comparison of results, it can be seen that brew with the highest total flavonoid content is from fresh fig leaf with a brewing temperature of 80°C. In addition, with the same brewing temperature, total flavonoid content of fresh fig leaf brew was higher than dried one. Based on previous research, the destruction of flavonoids had varied significant differences between fresh samples and dried materials. Heating may breakdown some phytochemicals which affect cell wall integrity and cause a migration of some flavonoids component. In addition, the loss in flavonoids may due to breakdown or leakage by chemical reactions includes oxygen, enzymes and light (Rababah *et al.*, 2015).

The data analysis with Two Way Anova test show that:

- a. brewing temperature significantly affects total flavonoid content,
- b. simplicia (fresh and dried) preparation significantly affects total flavonoid content, and
- c. the interactions of both (the brewing temperature and simplicia preparation) significantly affects total flavonoid content.

4. Conclusion

The total flavonoid contents of fresh fig leaf brew with the brewing temperatures of 60°C, 70°C, and 80°C are 0.0031±0.00039%, 0.0081±0.00021%, and 0.0092±0.00024%. The total flavonoid contents of dried fig leaf brew with the brewing temperatures of 60°C, 70°C, and 80°C are 0.0010±0.00025%, 0.0015±0.00004%, and 0.0017±0.00018%. Based on the results obtained, it can be concluded that increasing the brewing temperature from 60°C to 80°C increases the total flavonoid content of fig leaf brew, both fresh and dried. Total phenolic content of fresh fig leaf brew is higher than dried fig leaf brew, because the drying process decreases the availability of flavonoids. Brewing temperature, simplicia (fresh and dried) preparation, and the interactions of both (the brewing temperature and simplicia preparation) significantly affects total flavonoid content. For further research, it is necessary to increase the brewing temperature to find out the temperature when total flavonoid content decrease.

5. Acknowledgements

This research was funded by the Directorate of Research and Community Service, General Directorate of Research and Development Reinforcement, Ministry of Research, Technology and Higher Education of Republic of Indonesia with the novice lecturer research scheme 2018.

References

- Chandra, S., Khan, S., Avula, B., Lata, H., Yang, M. H., ElSohly, M. A., and Khan, I. A. (2014). Assessment of Total Phenolic and Flavonoid Content, Antioxidant Properties, and Yield of Aeroponically and Conventionally Grown Leafy Vegetables and Fruit Crops: A Comparative Study. *Evidence-Based Complementary and Alternative Medicine*. 2014: 1–9. <https://doi.org/10.1155/2014/253875>
- Jain, T., Jain, V., Pandey, R., Vyas, A., and Shukla, S. S. (2009). Microwave assisted extraction for phytoconstituents - an overview. *Asian Journal of Research in Chemistry*. 2(1): 19–25.
- Kozłowska, A. and Dorota, S.W. (2014). Flavonoids - Food Sources and Health Benefits. *Roczniki Państwowego Zakładu Higieny*. 65(2): 79–85.
- Kumar, R., Vijayalakshmi, S., and Nadanasabapathi, S. (2017). Health Benefits of Quercetin. *Defence Life Science Journal*. 2(2):142-148. <https://doi.org/10.14429/dlsj.2.11359>
- Mawa, S., Husain, K., and Jantan, I. (2013). *Ficus carica* L. (Moraceae): Phytochemistry, Traditional Uses and Biological Activities. *Evidence-Based Complementary and Alternative Medicine*. 2013: 1–8. <https://doi.org/10.1155/2013/974256>
- Pollastri, S. and Tattini, M. (2011). Flavonols: old compounds for old roles. *Annals of Botany*. 108(7): 1225–1233. <https://doi.org/10.1093/aob/mcr234>
- Rababah, T. M., Al-u'datt, M., Alhamad, M., Al-Mahasneh, M., Ereifej, K., Andrade, J., and Yang, W. (2015). Effects of drying process on total phenolics, antioxidant activity and flavonoid contents of common Mediterranean herbs. *International Journal of Agricultural and Biological Engineering*. 8(2): 145–150.
- Trifunski, S., Munteanu, M., Ardelean, D., Orodan, M., Osser, G., & Gligor, R. (2015). Flavonoids and polyphenols content and antioxidant activity of *Ficus carica* L. extracts from Romania. *Zbornik Matice Srpske Za Prirodne Nauke*, 32(128): 57–65. <https://doi.org/10.2298/ZMSPN1528057T>
- Vaya, J., and Mahmood, S. (2006). Flavonoid content in leaf extracts of the fig (*Ficus carica* L.), carob (*Ceratonia siliqua* L.) and pistachio (*Pistacia lentiscus* L.). *BioFactors*. 28(3–4): 169–175. <https://doi.org/10.1002/biof.5520280303>